maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to ompleting and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding an DMB control number.	ion of information. Send comments arters Services, Directorate for Info	regarding this burden estimate rmation Operations and Reports	or any other aspect of the property of the contract of the con	nis collection of information, Highway, Suite 1204, Arlington	
1. REPORT DATE 2005	2 DEDORT TVDE			3. DATES COVERED 00-00-2005 to 00-00-2005		
4. TITLE AND SUBTITLE	5a. CONTRACT NUMBER					
First Record of Culex (Culex) Coronator in Louisiana, USA				5b. GRANT NUMBER		
				5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Walter Reed Army Institute of Research, Department of Entomology, Silver Spring, MD, 20910				8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAII Approved for publ	LABILITY STATEMENT ic release; distributi	on unlimited				
13. SUPPLEMENTARY NO	OTES					
14. ABSTRACT see report						
15. SUBJECT TERMS						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	3		

Report Documentation Page

Form Approved OMB No. 0704-0188

SCIENTIFIC NOTE

FIRST RECORD OF *CULEX* (*CULEX*) *CORONATOR* IN LOUISIANA, USA

MUSTAPHA DEBBOUN, 1,3 DENNIS D. KUHR, 1 LEOPOLDO M. RUEDA2 AND JAMES E. PECOR2

ABSTRACT. The 1st confirmed record of *Culex coronator* for Louisiana was made at Fort Polk, LA, from carbon dioxide—baited light trap and gravid trap collections performed from April to October 2004. In addition to the new record, 17 mosquito species in 5 genera (*Aedes, Anopheles, Coquillettidia, Culex,* and *Psorophora*) were collected. Collection-site and species distribution data are included.

KEY WORDS Culex coronator, Louisiana, mosquitoes, Culicidae, Diptera

We report a new state record of mosquito species, *Culex* (*Culex*) *coronator* Dyar and Knab from Fort Polk, LA. This is one of 18 mosquito species collected from Centers for Disease Control and Prevention (CDC) light traps baited with carbon dioxide and a CDC gravid trap (John Hock Co., Gainesville, FL), from various sites in Fort Polk from April to October 2004.

Culex coronator has a wide geographic distribution, including Central and South America (Argentina, Belize, Bolivia, Brazil, Colombia, Costa Rica, El Salvador, French Guiana, Guatemala, Honduras, Nicaragua, Panama, Paraguay, Peru, Suriname, Trinidad and Tobago, and Venezuela) and the USA (Arizona, New Mexico, and Texas) (Darsie and Ward 2004, Walter Reed Bioystematics Unit 2001). Carpenter (1970) discounted the report of S. O. Hill, B. J. Smittle, and F. M. Philips in 1958 about the presence of this species in Louisiana (reference not available but cited by Carpenter [1970]). Our paper confirms the existence of this species in that state.

This species colonizes a wide variety of container, ground pool, and stream-associated habitats throughout its range. It has been collected in permanent and temporary sources of water, in the shade or in the sun, and in sylvatic and domestic habitats (Carpenter and LaCasse 1955; Horsfall 1972; Pecor et al. 2002; Debboun, unpublished data). On April 29, 2004, the Entomological Sciences Division (ESD) of the U.S. Army Center for Health Promotion and Preventive Medicine—South (USACHPPM-South) received the adult specimens from Fort Polk, LA, as part of the USACHPPM-South West Nile Virus Surveillance Program. The 1st specimens were collected on April 21, 2004, at

Twenty-three female specimens of Cx. coronator were collected in 2004, with 22 females from the CDC light traps and 1 from the CDC gravid trap (site no. 45, 31°02.29'N, 93°13.90'E). The descriptions of the trap sites in Fort Polk are as follows, with the date of collection and number of female Cx. coronator collected given in parentheses. Trap site no. 3 (31°02.08'N, 93°12.79'E, August 26, 2004, 10 females) was inside a tree line, across the street from the barracks, building no. 2300 and adjacent to an old beaver pond on Mississippi Avenue. Trap site no. 37 (31°02.42′N, 93°13.48′E, May 19, 2004, 1 female and on August 24, 2004, 7 females) was in the backyard of a housing quarters (no. 5116) overlooking a primarily dry, low area with a manhole cover approximately 30 feet away.

trap site no. 43 in Fort Polk (Fig. 1) by the U.S. Army Preventive Medicine Services personnel using a CDC light trap baited with carbon dioxide and sent to USACHPPM-South for identification and assay processing. These specimens were then sent to the Walter Reed Biosystematics Unit, Smithsonian Institution, for confirmation and were identified by J.E.P. and L.M.R. as Cx. coronator (Bram 1967, Darsie and Ward 2004). The classification used in this paper follows that of Knight and Stone (1977). Eight additional collections of Cx. coronator were made during 2004. A total of 18 species in 5 genera were collected from May to October 2004. The collected associated species were Aedes (Stegomyia) albopictus (Skuse), Anopheles (Anopheles) crucians Wiedemann, An. (Ano.) punctipennis (Say), An. (Ano.) quadrimaculatus Say, Coquillettidia perturbans (Walker), Cx. (Melanoconion) erraticus (Dyar and Knab), Cx. (Cux.) nigripalpus Theobald, Cx. (Cux.) quinquefasciatus Say, Cx. (Cux.) salinarius Coquillett, Aedes (Ochlerotatus) atlanticus Dyar and Knab/tormentor Dyar and Knab, Ae. (Och.) fulvus pallens Ross, Ae. (Och.) infirmatus Dyar and Knab, Ae. (Och.) thibaulti Dyar and Knab, Ae. (Och.) canadensis canadensis (Theobald), Ae. (Och.) sticticus (Meigen), Psorophora (Janthinosoma) ferox (Von Humboldt), and Ps. (Grabhamia) columbiae (Dyar and Knab).

¹ U.S. Army Center for Health Promotion and Preventive Medicine–South, Fort McPherson, GA 30330-1075.

² Department of Entomology, Walter Reed Army Institute of Research, Silver Spring, MD 20910-7500.

³ Present address: Academy of Health Sciences, Attn: MCCS-HPN, U.S. Army Medical Department Center and School, Fort Sam Houston, TX 78234-6142.




Fig. 1. Map of Fort Polk, LA, showing the collection site of *Culex coronator* where it was 1st collected on April 21, 2004

Trap site no. 43 (31°02.90′N, 93°12.94′E, May 25, 2004, 1 female) was behind the Veterinary Clinic kennels (building no. 665), located on a bluff about 20 ft above the tree canopy and overlooking a heavily forested area. Trap site no. 45 (31°02.29'N, 93°13.91′E, August 24, 2004, 1 female) was located near a sentinel chicken flock, near Magnolia Drive and between Monroe Drive and Michael Drive. This site was shady and damp, near a lift station (building no. 5292) in an old forest area next to a recently cleared sewage line right-of-way. Trap site no. 50 (31°02.74′N, 93°13.36′E, May 19, 2004, 1 female) was located in Norris Loop, near the 4900 block lift station, housing area, recently cleared forest area in the middle of housing with poor drainage and heavy organic debris. Trap site no. 51 (31°01.45'N, 93°12.12'E, October 6, 2004, 1 female) was in the Honor Field, to the left of the stands near the tree line. Trap site no. 52 (31°02.49′N, 93°12.18′E, May 25, 2004, 1 female) was on Texas Avenue, across from building no. 3334, in a recently cleared tree line with heavy organic debris.

Adult females of *Cx. coronator* have been collected from human bait in Brazil (Rachou et al. 1958, Roberts and Hsi 1979), and in Peru (Debboun, unpublished data). However, Carpenter and LaCasse (1955) noted that the females apparently do not feed on humans in the USA. This species also has been collected from chicken and rabbit baits in Argentina (Almiron and Brewer 1995). Turell et al. (2000) reported that *Cx. coronator* is not susceptible to epizootic (IAB and IC) and enzootic (ID and IE) strains of Venezuelan equine encephalomyelitis virus (VEE) in Peru. However, Scherer et al. (1971) isolated VEE from *Cx. coronator* in southeastern Mexico. Anderson et al. (1957) and

Aitken et al. (1969) also reported St. Louis encephalitis virus isolation from wild-collected female *Cx. coronator* in Trinidad. It is 1 of the 60 species found positive for West Nile virus (WNV) in mosquito pools in the USA from 1999 to 2004 (Centers for Disease Control and Prevention 2005).

Sample specimens of *Cx. coronator* and other associated species from CDC light traps and gravid traps collected from Fort Polk sites from May through October were all processed by the ESD, USACHPPM-South, assayed, and found negative for WNV and eastern equine encephalitis virus.

We thank Lenoir Ross of U.S. Army Preventive Medicine Services at Fort Polk for setting up the traps, collecting the adult mosquito specimens, and providing trap site descriptions, and Erin Stanwix, Jeffri Humphries, and Laura Frick for initial identification and laboratory assistance. Special thanks to Daniel Strickman and Richard Wilkerson for reviewing the manuscript and providing suggestions and advice. The opinions and assertions contained herein are those of the authors and are not to be construed as official or reflecting the views of the Department of Defense.

REFERENCES CITED

Aitken THG, Spence L, Jonkers AH, Downs WG. 1969.
A 10-year survey of Trinidadian arthropods for natural virus infections (1953–1963). J Med Entomol 6:207–215.

Almiron WR, Brewer MM. 1995. Host preference of Culicidae (Diptera) collected in central Argentina. *Rev Saude Publica* 29:108–114. [In Spanish.]

Anderson CR, Aitken THG, Downs WG, Spence L. 1957. The isolation of St. Louis virus from Trinidad mosquitoes. *Am J Trop Med Hyg* 6:688–692.

Bram RA. 1967. Classification of *Culex* subgenus *Culex* in the New World (Diptera: Culicidae). *Proc US Natl Mus Smithsonian Inst* 120:1–122.

Carpenter SL. 1970. Review of recent literature on mosquitoes of North America, supplement I. *Calif Vector Views* 17:39–66.

Carpenter SJ, LaCasse WJ. 1955. Mosquitoes of North America (north of Mexico) Berkeley, CA: Univ. Calif. Press

Centers for Disease Control and Prevention. 2005. West Nile virus. Division of Vector-Borne Infectious Diseases. National Center for Infectious Diseases, Fort Collins, CO. http://www.cdc.gov/ncidod/dvbid/westnile/mosquitospecies.htm

Darsie RF, Ward RA. 2004. *Identification and geographical distribution of the mosquitoes of North America, north of Mexico* Gainesville, FL: Univ. Press of Florida. Horsfall WR. 1972. *Mosquitoes: their bionomics and re-*

Horsfall WR. 1972. Mosquitoes: their bionomics and relation to disease New York: Hafner Publishing Co.

Knight, KL, Stone A. 1977. A catalog of the mosquitoes of the world (Diptera: Culicidae) Volume VI. College Park, MD: Thomas Say Foundation, Entomological Society of America.

Pecor, JP, Harbach RE, Peyton EL, Roberts DR, Rejmankova E, Manguin S, Palanko J. 2002. Mosquito studies in Belize, Central America: records, taxonomic notes, and a checklist of species. *J Am Mosq Control Assoc* 18:241–276.

- Rachou RG, Lima MM, Neto JAF, Martins CM. 1958. Algunas dados sobre o comportamento de mosquitos de Ponta Grossa (Florianopolis, Santa Catarina). Rev Bras Malariol Doencas Trop 9:417–427.
- Roberts DR, Hsi BP. 1979. An index of species abundance for use with mosquito surveillance data. *Environ Entomol* 8:1007–1013.
- Scherer WF, Dickerman RW, Diaz-Jajera A, Ward RA, Miller MH, Schaffer PA. 1971. Ecologic studies of Venezuelan encephalitis virus in southeastern Mexico. Am J Trop Med Hyg 20:969–979.
- Turell MJ, Jones JW, Sardelis MR, Dohm DJ, Coleman RE, Watts DM, Fernandez R, Calampa C, Klein TA. 2000. Vector competence of Peruvian mosquitoes (Diptera: Culicidae) for epizootic and enzootic strains of Venezuelan equine encephalomyelitis virus. *J Med En*tomol 37:835–839.
- Walter Reed Biosystematics Unit. 2001. Systematic catalog of Culicidae. Walter Reed Biosystematics Unit, Smithsonian Institution. Washington, DC. http://wrbu.si.edu/wrbu.html